




CONCLUSIONS

1. We found the Bethe Ansatz equations are the equations describing the vacuum configurations of certain quiver gauge theories in two dimensions
2. The duality to the spin chain requires certain relations between the masses of the matter fields to be obeyed. These masses follow naturally from the possibility to turn on the quasihomogeneous superpotentials (conformal fixed points)



Supersymmetric Quantum Field and String Theories and Integrable Lattice Models

Nikita Nekrasov
Landau-100
conference
Chernogolovka
June 25, 2008



Based on

NN, S. Shatashvili,
hep-th/0806...xyz

The Characters of our play

2,3, and 4 dimensional
susy gauge theories

With 4 supersymmetries
($\mathcal{N}=1$ $d=4$)

on the one hand

and

Quantum integrable systems
soluble by Bethe Ansatz

on the other hand



For example, we shall
relate
the XXX Heisenberg magnet
to
2d $N=2$ SYM theory
with some matter

Dictionary

$U(N)$ gauge theory with
 $N=4$ susy in two dimensions
with L fundamental hypermultiplets
softly broken to $N=2$ by giving
the generic twisted masses to the
adjoint, fundamental and antifundamental
chiral multiplets compatible with the
Superpotential inherited from the
 $N=4$ theory



Dictionary

Eigenstates of the spin chain Hamiltonian(s) are in one-to-one correspondence with the supersymmetric vacua of the gauge theory





Dictionary

*Eigenvalues of the spin chain Hamiltonians
coincide with the
vacuum expectation values
of the chiral ring operators
of the susy gauge theory*



Dictionary

$$H_k \Psi_\lambda = E_k(\lambda) \Psi_\lambda \iff E_k(\lambda) = \langle \lambda | \mathcal{O}_k | \lambda \rangle$$



Reminder on

$N=2$ supersymmetry in two dimensions: Basic multiplets

Vector multiplet contains a gauge field and adjoint complex scalar;

Chiral multiplet contains a (charged) complex scalar

(plus auxiliary bosonic field);

Twisted chiral multiplet contains a complex scalar and a gauge field strength;

Reminder on

$N=2$ supersymmetry in two dimensions: Basic Multiplets

Vector multiplet contains a gauge field and a complex scalar

$$V = \theta^- \bar{\theta}^- (v_0 - v_1) + \theta^+ \bar{\theta}^+ (v_0 + v_1) - \sqrt{2} \sigma \theta^- \bar{\theta}^+ - \sqrt{2} \bar{\sigma} \theta^+ \bar{\theta}^- \\ + 2i \theta^- \theta^+ (\bar{\theta}^- \bar{\lambda}_- + \bar{\theta}^+ \bar{\lambda}_+) + 2i \bar{\theta}^+ \bar{\theta}^- (\theta^+ \lambda_+ + \theta^- \lambda_-) + 2 \theta^- \theta^+ \bar{\theta}^+ \bar{\theta}^- D.$$

Reminder on $\mathcal{N}=2$ supersymmetry in two dimensions: Basic Multiplets

Chiral multiplet contains a complex scalar
(plus auxiliary bosonic field);

$$\Phi = \phi + \sqrt{2}\theta^\alpha\psi_\alpha + \theta^\alpha\theta_\alpha F.$$

Reminder on $N=2$ supersymmetry in two dimensions: Basic Multiplets

Chiral multiplet contains a complex scalar
(plus auxiliary bosonic field);

$$\bar{D}_\alpha \Phi = 0.$$

Reminder on
 $N=2$ supersymmetry in two
dimensions: Basic Multiplets

Chiral multiplets will be denoted by

Q , \tilde{Q} , and Φ .

Reminder on N=2 supersymmetry in two dimensions: Basic Multiplets

Twisted chiral multiplet contains a gauge field strength and a complex scalar

$$\Sigma = \frac{1}{2\sqrt{2}} \{ \bar{\mathcal{D}}_+, \mathcal{D}_- \} = \sigma + i\sqrt{2}\theta^+ \bar{\lambda}_+ - i\sqrt{2}\bar{\theta}^- \lambda_- + \sqrt{2}\theta^+ \bar{\theta}^- D$$

..... $- i\sqrt{2}\theta^+ \bar{\theta}^- F_{01}$

$$\bar{\mathcal{D}}_+ \Sigma = \mathcal{D}_- \Sigma = 0.$$

Reminder on

$N=2$ supersymmetry in two dimensions: Basic Multiplets

Twisted chiral multiplet contains a gauge field strength and a complex scalar:

Full expansion

$$\begin{aligned}\Sigma = \frac{1}{2\sqrt{2}}\{\bar{\mathcal{D}}_+, \mathcal{D}_-\} &= \sigma + i\sqrt{2}\theta^+\bar{\lambda}_+ - i\sqrt{2}\bar{\theta}^-\lambda_- + \sqrt{2}\theta^+\bar{\theta}^- D \\ &- i\bar{\theta}^-\theta^-(D_0 - D_1)\sigma - i\theta^+\bar{\theta}^+(D_0 + D_1)\sigma \\ &+ \sqrt{2}\bar{\theta}^-\theta^-\theta^+(D_0 - D_1)\bar{\lambda}_+ - \sqrt{2}\theta^+\bar{\theta}^+\bar{\theta}^-(D_0 + D_1)\lambda_- - i\sqrt{2}\theta^+\bar{\theta}^- F_{01} \\ &- 2i\bar{\theta}^-\theta^-\theta^+[\sigma, \bar{\lambda}_+] - 2i\bar{\theta}^-\theta^+\bar{\theta}^+[\sigma, \lambda_-] \\ &- \bar{\theta}^-\theta^-\theta^+\bar{\theta}^+ ((D_0^2 - D_1^2)\sigma - [\sigma, [\sigma, \bar{\sigma}]] + i\bar{\theta}^-\theta^-\theta^+\bar{\theta}^+[\sigma, \partial_m v^m]).\end{aligned}$$

Reminder on $\mathcal{N}=2$ supersymmetry in two dimensions: Lagrangians

Gauge kinetic terms

$$\begin{aligned} L_g &= -\frac{1}{4e^2} \int d^2x d^4\theta \operatorname{Tr} \bar{\Sigma} \Sigma \\ &= \frac{1}{e^2} \int d^2x \operatorname{Tr} \left(\frac{1}{2} F_{01}^2 + |D_0 \sigma|^2 - |D_1 \sigma|^2 + i \bar{\lambda}_- (D_0 + D_1) \lambda_- + i \bar{\lambda}_+ (D_0 - D_1) \lambda_+ \right. \\ &\quad \left. + \frac{1}{2} D^2 - \frac{1}{2} [\sigma, \bar{\sigma}]^2 - \sqrt{2} \lambda_+ [\sigma, \bar{\lambda}_-] + \sqrt{2} [\bar{\sigma}, \lambda_-] \bar{\lambda}_+ \right). \end{aligned}$$

*Reminder on
N=2 supersymmetry in two
dimensions: Lagrangians*

Fayet-Iliopoulos and theta terms

$$\begin{aligned}\mathcal{L}_{\text{FI},\theta} &= \frac{i\tau}{4} \int d^2\tilde{\theta} \text{Tr}\Sigma + h.c. \\ &= \int d^2x \left(-r \text{Tr} D + \frac{\theta}{2\pi} \text{Tr} F_{01} \right)\end{aligned}$$

*Reminder on
N=2 supersymmetry in two
dimensions: Lagrangians*

Fayet-Iliopoulos and theta terms

$$\mathcal{L}_{\text{FI},\theta} = \frac{i\tau}{4} \int d^2\tilde{\theta} \text{Tr}\Sigma + h.c.$$

Give an example of the twisted superpotential

$$\tau = ir + \theta/2\pi.$$

*Reminder on
N=2 supersymmetry in two
dimensions: Lagrangians*

Matter kinetic terms

$$L_{ch} = \frac{1}{4} \int d^2x d^4\theta \bar{\Phi}\Phi$$

Reminder on $\mathcal{N}=2$ supersymmetry in two dimensions: Lagrangians

Matter kinetic terms

$$\begin{aligned} &= \int d^2x \left(|D_0\phi|^2 - |D_1\phi|^2 + |F|^2 + i\bar{\psi}_+(D_0 - D_1)\psi_+ \right. \\ &\quad + i\bar{\psi}_-(D_0 + D_1)\psi_- + \bar{\phi}D\phi - \bar{\phi}\{\sigma, \bar{\sigma}\}\phi \\ &\quad - \sqrt{2}\bar{\psi}_+\bar{\sigma}\psi_- - \sqrt{2}\bar{\psi}_-\sigma\psi_+ + i\sqrt{2}\bar{\psi}_+\bar{\lambda}_-\phi - i\sqrt{2}\bar{\psi}_-\bar{\lambda}_+\phi \\ &\quad \left. + i\sqrt{2}\bar{\phi}\lambda_+\psi_- - i\sqrt{2}\bar{\phi}\lambda_-\psi_+ \right). \end{aligned}$$

*Reminder on
N=2 supersymmetry in two
dimensions: Lagrangians*

Superpotential terms

$$L_W = - \int d^2y d\theta^+ d\theta^- W(\Phi_i)|_{\bar{\theta}^+ = \bar{\theta}^- = 0} - h.c.$$

*Reminder on
N=2 supersymmetry in two
dimensions: Lagrangians*

Superpotential terms

$$L_W = - \int d^2y \left(F_i \frac{\partial W}{\partial \phi_i} + \frac{\partial^2 W}{\partial \phi_i \partial \phi_j} \psi_{-,i} \psi_{+,j} \right) - h.c.$$

*Reminder on
N=2 supersymmetry in two
dimensions: Lagrangians*

Twisted superpotential terms

$$\Delta L = \int d^2y d\theta^+ d\bar{\theta}^- \widetilde{W}(\Sigma) \Big|_{\theta^- = \bar{\theta}^+ = 0} + h.c.$$

*Reminder on
N=2 supersymmetry in two
dimensions: Lagrangians*

Twisted superpotential terms

$$= \int d^2y \left(\sqrt{2} \widetilde{W}'(\sigma) (D - i\nu_{01}) + 2 \widetilde{W}''(\sigma) \bar{\lambda}_+ \lambda_- \right) + h.c.$$

Reminder on
 $N=2$ supersymmetry in two
dimensions: Lagrangians

Twisted mass terms

$$\mathcal{L}_{\text{mass}} = \int d^4\theta \operatorname{tr} \left(\Phi^\dagger e^{\tilde{V}} \otimes \operatorname{Id}_{\text{color space}} \Phi \right)$$

Reminder on $\mathcal{N}=2$ supersymmetry in two dimensions: Lagrangians

Twisted mass terms

$$\mathcal{L}_{\text{mass}} = \int d^4\theta \operatorname{tr} \left(\Phi^\dagger e^{\tilde{V}} \otimes \operatorname{Id}_{\text{color space}} \Phi \right)$$

where $\tilde{V} = \tilde{m} \theta_+ \bar{\theta}_-$, \tilde{m} acts in a flavour space, and, to preserve susy:

$$[\tilde{m}, \tilde{m}^*] = 0$$

Reminder on $N=2$ supersymmetry in two dimensions: Lagrangians

Twisted mass terms

$$\mathcal{L}_{\text{mass}} = \int d^4\theta \operatorname{tr} \left(\Phi^\dagger e^{\tilde{V}} \otimes \operatorname{Id}_{\text{color space}} \Phi \right)$$

$$\tilde{V} = \tilde{m} \theta_+ \bar{\theta}_- \quad - \text{Background vector field for global symmetry}$$

Reminder on $N=2$ supersymmetry in two dimensions: Lagrangians

Cf. the ordinary mass terms

$$\mathcal{L}_{\text{mass}} = \sum_{i, \tilde{j}} \int d^2\theta \, m_i^{\tilde{j}} \tilde{Q}_{\tilde{j}} Q^i + \text{h.c.},$$

Which are just the superpotential terms

General strategy

Take an $\mathcal{N}=2$ $d=2$ gauge theory with matter,
in some representations \mathbf{R}_f
of the gauge group \mathbf{G}
integrate out the massive matter fields,
compute
the effective twisted super-potential
on the Coulomb branch

$$W_{\text{eff}} = \sum_f \text{Tr}_{\mathbf{R}_f} (\sigma + m_f) (\log (\sigma + m_f) - 1) + 2\pi i \tau \text{Tr} \sigma$$

Vacua of the gauge theory

For $G = U(N)$

$$\sigma \rightarrow \text{diag}(\sigma_1, \dots, \sigma_N)$$

$$\frac{\partial W_{\text{eff}}}{\partial \sigma_i} = \lambda_i - i + \frac{1}{2}(N + 1)$$

Due to quantization of the gauge flux

$$\lambda_i \in \mathbf{Z}$$

Familiar example:

CP^N model

Field content:

(N+1) chiral multiplet of charge +1

Q^i $i=1, \dots, N+1$

U(1) gauge group

σ is a scalar

Familiar example:

CP^N *model*

Effective twisted superpotential

(D'ADDA, A.LUSCHER, DI VECCHIA)

$$(\mathbf{N} + 1)\sigma(\log\sigma - 1) + 2\pi i\tau \sigma$$

N+1 vacuum

$$\sigma^{\mathbf{N}+1} = e^{2\pi i\tau}$$

**Quantum
cohomology**

More interesting example

Field content

Gauge group: $G=U(N)$ $\sigma \rightarrow \text{diag}(\sigma_1, \dots, \sigma_N)$

Matter chiral multiplets:

1 <u>Adjoint</u>	twisted mass	M
N_f <u>fundamentals</u>	... mass	m_f
N_f <u>anti-fundamentals</u>	... mass	$m_{\bar{f}}$

More interesting example

Effective superpotential:

$$\begin{aligned} & N_f (\sigma_i + m_f) (\log (\sigma_i + m_f) - 1) + \\ & N_{\bar{f}} (-\sigma_i + m_{\bar{f}}) (\log (-\sigma_i + m_{\bar{f}}) - 1) + \\ & \sum_{i,j} (\sigma_i - \sigma_j + M) (\log (\sigma_i - \sigma_j + M) - 1) \\ & + (N_{\bar{f}} - N_f) \log \Lambda \sum_i \sigma_i \end{aligned}$$

More interesting example

Equations for vacua:

$$\Lambda^{N_f - N_f} \frac{\left(\sigma_i + m_f\right)^{N_f}}{\left(-\sigma_i + m_{\bar{f}}\right)^{N_f}} = \prod_{j \neq i} \frac{\sigma_i - \sigma_j + M}{\sigma_i - \sigma_j + M}$$

More interesting example

Non-anomalous, UN finite case.

$$N_f = N_{\bar{f}} = L$$

More interesting example

Non-anomalous, UV finite case.

$$N_f = N_{\bar{f}} = L$$

Redefine:

$$\sigma_j = \frac{1}{2} (m_{\bar{f}} - m_f) - iM\lambda_j$$

$$\frac{1}{2} (m_{\bar{f}} + m_f) = Ms$$

Vacua of gauge theory

$$\left(\frac{\lambda_i + iS}{\lambda_i - iS} \right)^L = \prod_{j \neq i} \frac{\lambda_i - \lambda_j + i}{\lambda_i - \lambda_j - i}$$

Vacua of gauge theory


$$\left(\frac{\lambda_i + iS}{\lambda_i - iS} \right)^L = \prod_{j \neq i} \frac{\lambda_i - \lambda_j + i}{\lambda_i - \lambda_j - i} \times e^t$$

$$t = r + i\vartheta$$

Gauge theory - spin chain

Identical
to the
Bethe
equations
for spin s
XXX magnet


$$\left(\frac{\lambda_i + is}{\lambda_i - is} \right)^L = \prod_{j \neq i} \frac{\lambda_i - \lambda_j + i}{\lambda_i - \lambda_j - i}$$



Gauge theory - spin chain

Identical to
the Bethe
equations for
spin s XXX
magnet
with twisted
boundary
conditions

$$\left(\frac{\lambda_i + is}{\lambda_i - is} \right)^L = \prod_{j \neq i} \frac{\lambda_i - \lambda_j + i}{\lambda_i - \lambda_j - i} \times e^t$$



Gauge theory - spin chain

Gauge theory
vacua -
eigenstates of the
spin Hamiltonian
(transfer-matrix)

$$H = \sum_{n=1}^L \sigma_n^a \otimes \sigma_{n+1}^a$$




Table of dualities

XXX spin chain

$SU(2)$

L spins

N excitations

$U(N)$ $d=2$ $N=2$

Chiral

multiplets:

1 adjoint

L fundamentals

L anti-fund.

$$\left(\frac{\lambda_i + is}{\lambda_i - is}\right)^L = \prod_{j \neq i} \frac{\lambda_i - \lambda_j + i}{\lambda_i - \lambda_j - i}$$

NB: Special masses, to be explained

Table of dualities

XXZ spin chain

$SU(2)$

L spins

N excitations

$U(N)$ $d=3$ $N=2$

Compactified on a circle

Chiral multiplets:

1 adjoint

L fundamentals

L anti-fund.

$$\left(\frac{\sinh(\lambda_i + i\gamma)}{\sinh(\lambda_i - i\gamma)} \right)^L = \prod_{j \neq i} \frac{\sinh(\lambda_i - \lambda_j + i\gamma)}{\sinh(\lambda_i - \lambda_j - i\gamma)}$$

Special masses again

Table of dualities

XYZ spin chain

$SU(2)$, $L = 2N$

spins

N excitations

$$H = \sum_{n=1}^L J_x \sigma_n^x \otimes \sigma_{n+1}^x + J_y \sigma_n^y \otimes \sigma_{n+1}^y + J_z \sigma_n^z \otimes \sigma_{n+1}^z$$

$U(N)$ $d=4$ $N=1$

Compactified on a

2-torus = elliptic curve E

Chiral multiplets:

1 adjoint

$L = 2N$ fundamentals

$L = 2N$ anti-fund.

**Masses = wilson loops
of the flavour group**

Table of dualities

XYZ spin chain

$SU(2)$, $L = 2N$ spins

N excitations

$$H = \sum_{n=1}^L J_x \sigma_n^x \otimes \sigma_{n+1}^x + J_y \sigma_n^y \otimes \sigma_{n+1}^y + J_z \sigma_n^z \otimes \sigma_{n+1}^z$$

$U(N)$ $d=4$ $N=2$

Compactified on a

2-torus = elliptic curve E

$L = 2N$ fundamental
hypermultiplets

**Softly broken down to $N=1$ by the
wilson loops of the global
symmetry group = flavour group
 $U(L) \times U(1)$
= points on the Jacobian of E**

Table of dualities

It is remarkable that the spin chain has precisely those generalizations: rational (XXX), trigonometric (XXZ) and elliptic (XYZ) that can be matched to the 2, 3, and 4 dim cases.

$$H = \sum_{n=1}^L J_x \sigma_n^x \otimes \sigma_{n+1}^x + J_y \sigma_n^y \otimes \sigma_{n+1}^y + J_z \sigma_n^z \otimes \sigma_{n+1}^z$$

$$J_x = 1 + k \operatorname{sn}^2 2\eta, \quad J_y = 1 - k \operatorname{sn}^2 2\eta, \quad J_z = \operatorname{cn} 2\eta \operatorname{dn} 2\eta$$

Table of dualities

The L fundamentals and L anti-fundamentals can have different twisted masses

Field	Twisted mass
\tilde{Q}_a	$\mu_a + i s_a M$
Q^a	$-\mu_a + i s_a M$
Φ	$-iM$

This theory maps to inhomogeneous spin chain with different spins at different sites

Table of dualities

Yang-Yang counting function =
effective twisted superpotential



Table of dualities

Commuting hamiltonians (expansion of transfer matrix) =
the chiral ring generators, like

$$\text{Tr} \sigma^M$$

Table of dualities

Gauge theory theta angle (complexified)
is mapped to the spin chain theta angle
(twisted boundary conditions)



Algebraic Bethe Ansatz

Faddeev et al.

The spin chain is solved algebraically
using certain operators,

$$A(\lambda), B(\lambda), C(\lambda), D(\lambda)$$

obeying exchange commutation
relations


Algebraic Bethe Ansatz

The eigenvectors, Bethe vectors, are obtained by applying these operators to the (pseudo)vacuum.

$$\Psi_{\vec{\lambda}} = B(\lambda_1)B(\lambda_2)\dots B(\lambda_N)\Omega$$

Algebraic Bethe Ansatz VS GAUGE THEORY

For the spin chain it is natural to fix $L =$
total number of spins
and consider various $N =$ excitation levels



In the gauge theory context N is fixed.

Algebraic Bethe Ansatz VS STRING THEORY

However, if the theory is embedded
into string theory via brane
realization

then changing N is easy:
bring in an extra brane.

One might use the constructions
of Witten'96, Hanany-Hori'02

Algebraic Bethe Ansatz VS STRING THEORY

THUS:

$B(\lambda)$ is for BRANE!

λ is for location!

Are these models
too special, or the
gauge theory/integrable
lattice model
correspondence is
more general?

Actually, virtually any
Bethe ansatz soluble
system can be mapped to a
 $N=2$ $d=2$ gauge theory:
General spin group \mathcal{H} ,
8-vertex model,
Hubbard model,

More general spin chains

The $SU(2)$ spin chain
has generalizations to
other groups and representations.

Quoting the (nested) Bethe ansatz
equations from N. Reshetikhin

General groups/ reps

For simply-laced group \mathcal{H} of rank r

$$N \longrightarrow \sum_{\mathbf{i}=1}^r N_{\mathbf{i}}, \quad L \longrightarrow \sum_{\mathbf{i}=1}^r L_{\mathbf{i}}$$

$$i \longrightarrow (\alpha, \mathbf{i}); \quad \mathbf{i} = 1, \dots, r, \quad \alpha = 1, \dots, N_{\mathbf{i}}$$

General groups/ reps


For simply-laced group \mathcal{H} of rank r

$$\prod_{a=1}^{L_i} \frac{\lambda_{\alpha}^{(i)} - \mu_a^{(i)} + s_a^{(i)}}{\lambda_{\alpha}^{(i)} - \mu_a^{(i)} - s_a^{(i)}} = \prod_{j=1}^r \prod_{\beta} \frac{\lambda_{\alpha}^{(i)} - \lambda_{\beta}^{(j)} + C_{ij}}{\lambda_{\alpha}^{(i)} - \lambda_{\beta}^{(j)} - C_{ij}}$$

(μ_a^i, s_a^i) Label representations of the Yangian of \mathcal{H} : Kirillov-Reshetikhin modules


C_{ij} Cartan matrix of \mathcal{H}

General groups/ reps
from GAUGE THEORY



QUIVER GAUGE THEORY

Symmetries




gauge group: $U(N_1) \times U(N_2) \times \dots \times U(N_r)$

flavor group: $U(L_1)^2 \times U(L_2)^2 \times \dots \times U(L_r)^2$

QUIVER GAUGE THEORY

Symmetries



gauge group: $U(N_1) \times U(N_2) \times \dots \times U(N_r)$

flavor group: $U(L_1)^2 \times U(L_2)^2 \times \dots \times U(L_r)^2$

QUIVER GAUGE THEORY

Charged matter



Adjoint chiral multiplet



Fundamental chiral multiplet




Anti-fundamental chiral multiplet



Bi-fundamental chiral multiplet


QUIVER GAUGE THEORY

Matter fields: adjoints




QUIVER GAUGE THEORY

Matter fields: fundamentals +
antifundamentals




QUIVER GAUGE THEORY

Matter fields: bi-fundamentals



QUIVER GAUGE THEORY


Full assembly in the $N=2$ $d=2$ language



QUIVER GAUGE THEORY: twisted masses

Adjoints


MC_{ii}



QUIVER GAUGE THEORY: twisted masses

fundamentals
anti-fundamentals

$$M(\mu_a^{(i)} \pm s_a^{(i)})$$




$$a = 1, \dots, L_i$$

QUIVER GAUGE THEORY: twisted masses

Bi-fundamentals

MC_{ij}



What is so special
about all these
masses?

The twisted masses
correspond to
symmetries.

Symmetries are
restricted by the e.g.
superpotential
deformations

The $N=2$ $d=4$ ($N=4$ $d=2$)
superpotential

$$W_0 = \sum_{a=1}^L \text{tr } \tilde{Q}_a \Phi Q^a$$

Has a symmetry

$$Q^a \mapsto e^{i\beta_a - i\frac{1}{2}\gamma} Q^a, \quad \tilde{Q}_b \mapsto e^{-i\beta_b - i\frac{1}{2}\gamma}, \quad \Phi \mapsto e^{i\gamma} \Phi$$

$$W_0 = \sum_{a=1}^L \text{tr} \tilde{Q}_a \Phi Q^a$$

It is this symmetry

$$Q^a \mapsto e^{i\beta_a - i\frac{1}{2}\gamma} Q^a, \quad \tilde{Q}_b \mapsto e^{-i\beta_b - i\frac{1}{2}\gamma}, \quad \Phi \mapsto e^{i\gamma} \Phi$$

*which
explains the ratio of
adjoint and
Fundamental masses*

Similarly, we should ask:
Why choose s_a
Half-integral
in the table

Field	Twisted mass
\tilde{Q}_a	$\mu_a + i s_a M$
Q^a	$-\mu_a + i s_a M$
Φ	$-i M$

?

The answer is that one
can turn on more
general superpotential

(ONLY N=2 D=2 IS PRESERVED)

$$W_{\tilde{Q}\Phi Q} = \sum_{a=1}^L u_a \tilde{Q}_a \Phi^{2s_a} Q^a$$

Which has a symmetry

$$Q^a \mapsto e^{i\beta_a - is_a \gamma} Q^a, \tilde{Q}_b \mapsto e^{-i\beta_b - is_b \gamma}, \Phi \mapsto e^{i\gamma} \Phi$$

Hubbard model

Bethe ansatz

$$e^{ik_\alpha L} = \prod_{l=1}^M \frac{\lambda_l - \sin k_\alpha + iU/4}{\lambda_l - \sin k_\alpha - iU/4}, \quad j = 1, \dots, N$$
$$\prod_{\alpha=1}^N \frac{\lambda_l - \sin k_\alpha - iU/4}{\lambda_l - \sin k_\alpha + iU/4} = \prod_{m=1}^M \frac{\lambda_l - \lambda_m - iU/2}{\lambda_l - \lambda_m + iU/2}, \quad l = 1, \dots, M$$


Lieb-Wu

The new ingredient is the
coupling to
a nonlinear sigma model

Mirrors
to **CP¹**

$$\begin{aligned}
 -iW_{\vec{n}, \vec{m}}(\vec{k}, \vec{\mu}, \vec{\sigma}) = & \\
 & -i\ell \sum_{a=1}^L (k_a \mu_a + \cos k_a) \\
 & + \sum_{\alpha=1}^N \sum_{a=1}^L (\sigma_\alpha - \mu_a + u) [\log(\sigma_\alpha - \mu_a + u) - 1] + \\
 & + \sum_{\alpha=1}^N \sum_{a=1}^L (\mu_a - \sigma_\alpha + u) [\log(\mu_a - \sigma_\alpha + u) - 1] \\
 & + \sum_{\alpha, \beta=1}^N (\sigma_\alpha - \sigma_\beta - 2u) [\log(\sigma_\alpha - \sigma_\beta - 2u) - 1] \\
 & + 2\pi i \sum_{a=1}^L n_a \mu_a + 2\pi i \sum_{\alpha=1}^N m_\alpha \sigma_\alpha
 \end{aligned}$$

Finally, what is the meaning of the spins?



What is the meaning of
Bethe wavefunction?

If time permits....

Further
developments:
Instanton corrected
Bethe Ansatz
equations

Instanton corrected Bethe Ansatz equations

Consider

$\mathcal{N}=2^*$ theory on $\mathbf{R}^2 \times \mathbf{S}^2$

with a partial twist along the two-sphere

One gets a deformation of the

Yang-Mills-Hitchin theory

(introduced in Moore-NN-Shatashvili'97)

(if \mathbf{R}^2 is replaced by a Riemann surface)

Twisted superpotential from prepotential

$$\tilde{W}_{\text{eff}}(a; m, \tau, \lambda) = \frac{\partial \mathcal{F}}{\partial m} + \frac{1}{2\pi i} \frac{\partial \mathcal{F}}{\partial \tau} + \sum_i \left(n_i a^i + m^i a_{D,i} \right)$$

Induced by twist *Tree level part* *Flux superpotential*
(Losev-NN-Shatashvili'97)

Twisted superpotential from prepotential

Magnetic flux $m^i = \int_{S^2} F^i$

Electric flux $n_i = \lambda_i - i + \frac{1}{2}(N + 1)$

In the limit of vanishing S^2
the magnetic flux should vanish

Twisted superpotential from prepotential

The prepotential is known explicitly as a series in $q = e^{2\pi i\tau}$:

$$\mathcal{F} = \mathcal{F}^{\text{pert}} + q\mathcal{F}_1 + q^2\mathcal{F}_2 + \dots$$

$$\mathcal{F}^{\text{pert}} = \pi i\tau \sum_i a_i^2 + \frac{1}{2} \sum_{i \neq j} a_{ij}^2 \left(\log(a_{ij}) - \frac{3}{2} \right) - (a_{ij} + m)^2 \left(\log(a_{ij} + m) - \frac{3}{2} \right)$$

$$\mathcal{F}_1 = m^2 \sum_i T_i$$

$$\mathcal{F}_2 = \sum_i \left(-\frac{3m^2}{2} T_i^2 + \frac{m^4}{2} T_i T_i^{(2)} \right)$$

$$+ m^6 \sum_{i \neq j} \frac{T_i T_j}{a_{ij}^2} \frac{(m^2 - 3a_{ij}^2)}{(a_{ij}^2 - m^2)^2}$$

$$T_i(s) = \prod_{j \neq i} \left(1 - \frac{m^2}{(s + a_{ij})^2} \right) = T_i + \frac{s^2}{2} T_i^{(2)} + o(s^2)$$

*Instanton corrected
Bethe Ansatz
equations*

$$\exp \left\{ 2\pi i \frac{\partial^2 \mathcal{F}}{\partial a_i \partial m} + \sum_k t_k \frac{du_k}{da_i} \right\} = 1$$

*We can read off an «S-matrix»
It contains 2-, 3-, higher order
interactions*

Instanton corrected Bethe Ansatz equations

The prepotential of the low-energy effective theory
is governed by a classical (holomorphic) integrable system

Donagi-Witten'95

Liouville tori = Jacobians of Seiberg-Witten curves

Classical integrable system
vs
Quantum integrable system

That system is quantized when
the gauge theory is subject to
the Omega-background


NN'02
NN-Okounkov'03
Braverman'03

Our quantum system is different!

Blowing up the two-sphere

2d Hitchin configuration

4d instanton




Wall-crossing phenomena
(new states, new solutions)

Something for the future

Remark

In the $s_a^{(i)} \rightarrow 0$ limit
the supersymmetry enhances

Cf. BA for QCD
Lipatov, Faddeev-Korchemsky'94



CONCLUSIONS

3. The algebraic Bethe ansatz seems to provide a realization of the brane creation operators -- something of major importance both for topological and physical string theories
4. Obviously this is a beginning of a beautiful story....